

gb Basic PCR Master Mix

Description of the product

gb Basic PCR Master Mix consists of hot-start Taq DNA polymerase, reaction buffer, dNTP, MgCl₂ and additives that prevent PCR inhibition. Taq polymerase is chemically modified DNA polymerase from *Thermus aquaticus*. This polymerase is completely inactive at room temperature but is quickly activated during the initial denaturation step of PCR.

Purpose of the product

gb Basic PCR Master Mix is intended for research end-point PCR applications. Beneficial feature is its enhanced resistance to PCR inhibitors. It is not intended for use in diagnostics.

gb PCR Master Mix by application	gb Basic
end-point PCR, common PCR amplification	✓
real-time PCR without probes	
real-time PCR with hydrolysis probes	
real-time PCR with LNA probes	
real-time PCR with hybridization probes	✓
real-time PCR with High Resolution Melting Analysis	
real-time PCR with low DNA samples	
PCR/real-time PCR with inhibited samples	✓

Available products

Cat. No.	Product	rxn
3004	gb Basic PCR Master Mix	100

1 tube contains reagents to provide 100 PCR reactions (20 µl volume of each reaction).

Parameters of the product

- Master Mix is a **2× concentrated solution**.
- It contains **all the components** necessary for PCR performance.
- It is specified especially for real-time PCR applications demanding **high sensitivity**, for example microbial DNA detection and somatic mutations detection.
- **Polymerase is a hot-start type** with a short activation time (3 min / 95 °C), with 5'-3' polymerase and exonuclease activity, 3'-5' exonuclease activity is not present.

Amplification protocol

Step	Temperature	Time	Cycle number
Initial denaturation/enzyme activation	95 °C	1–2 min	1
Denaturation	95 °C	0.5–1 min	
Annealing	T _m - 5 °C	0.5–1 min	30 - 50
Extension	72 °C	1 min/kb	
Final extension	72 °C	5–15 min	1